

Routine childhood immunisations from June 2013

When to immunise	Diseases protected against	Vaccine given	Immunisation site**
Two months old	Diphtheria, tetanus, pertussis, polio and Haemophilus influenzae type b (Hib)	DTaP/IPV/Hib (PediaceL)	Thigh
	Withdrawn August 2015		
			Thigh
Three months old	Diphtheria, tetanus, pertussis, polio and Hib	DTaP/IPV/Hib (PediaceL)	Thigh
	Meningococcal group C disease (MenC)	Men C (NeisVac-C or Menjugate)	Thigh
	Rotavirus (from July)	Rotavirus (Rotarix)	By mouth
Four months old	Diphtheria, tetanus, pertussis, polio and Hib	DTaP/IPV/Hib (PediaceL)	Thigh
	Pneumococcal disease	PCV (Prevenar 13)	Thigh
Between 12 and 13 months old – within a month of the first birthday	Hib/MenC	Hib/MenC (Menitorix)	Upper arm/thigh
	Pneumococcal disease	PCV (Prevenar 13)	Upper arm/thigh
	Measles, mumps and rubella (German measles)	MMR (Priorix or MMR VaxPRO)	Upper arm/thigh
Three years four months old or soon after	Diphtheria, tetanus, pertussis and polio	dTaP/IPV (Repevax) or DTaP/IPV (Infanrix-IPV)	Upper arm
	Measles, mumps and rubella	MMR (Priorix or MMR VaxPRO) (check first dose has been given)	Upper arm
Girls aged 12 to 13 years old	Cervical cancer caused by human papillomavirus types 16 and 18 (and genital warts caused by types 6 and 11)	HPV (Gardasil)	Upper arm
Around 14 years old	Tetanus, diphtheria and polio	Td/IPV (Revaxis), and check MMR status	Upper arm
	MenC†	Men C (Meningitec, Menjugate or NeisVac-C)††	Upper arm

** Where two or more injections are required at once, these should ideally be given in different limbs. Where this is not possible, injections in the same limb should be given 2.5cm apart. For more details see Chapters 4 and 11 in the Green Book.

† This vaccination will be introduced during the 2013/14 academic year.

†† The vaccine supplied will depend on the brands available at the time of ordering.

Immunisations for at-risk children

At birth, 1 month old, 2 months old and 12 months old	Hepatitis B	Hep B	Thigh
At birth	Tuberculosis	BCG	Upper arm (intradermal)



*NB Where a vaccine is manufactured by more than one supplier, it may, on occasion be necessary to substitute an alternative brand.

All vaccines for use in the routine childhood programme are available free of charge at www.immform.dh.gov.uk